

THERMOMETRUM MAGNUM:
O R,
GRAND STANDARD
THERMOMETER.

EXPRESSING

All Degrees of HEAT and COLD, from that with
which Mercury BOILS, to that which congeals it
into *solid* METAL.

TO WHICH ARE ADJUSTED

The celebrated SCALES of SIR ISAAC NEWTON, FAHRENHEIT,
De L'ISLE, and REAUMUR, for comparing Observations
made in every Part of the Globe, and in all Degrees of
Temperature in the Air, or any other Bodies.

The Whole delineated on, and illustrated by, a Large
COPPER-PLATE.

BY BENJAMIN MARTIN.

L O N D O N:

Printed for, and sold by the AUTHOR,

In Fleet-street, No 171, and by the Bookfellers in City
and Country. A D. 1772.



MERCURY boils

LEAD melts

BISMUTH melts

TIN melts

WATER boils

WAX melts

BLOOD Heat

SUMMER Heat

TEMPERATE


FREEZING Point


FREEZING Mixture

AIR at Petersburg

Common Freez^e Mixture
at Peter sburgh


Strong Freez^e Mixture
at Petersburg





THERMOMETRUM
 MAGNUM
 or the
 STANDARD
 THERMOMETER

FAHRENHEIT
 DE LISSE
 NEWTON
 STANDARD
 REAUMUR




A

DESCRIPTION

OF THE STANDARD

THERMOMETER.

 S I have but slightly touched upon the Nature and Use of the THERMOMETER in the Treatise on the BAROMETER, it may be proper here to give a further Account of an Instrument so generally useful, in itself so very curious, and which has arrived to it's greatest Perfection only within twelve Years past, by the singular and amazing Invention of Mr. J. A. BRAUNIVS, of the *Imperial* ACADEMY OF SCIENCES at PETERSBURGH ; of which the following is an Abstract, translated from his own Dissertation on the Subject

The *Thermometer* he used for the Purposes of exploring the most wonderful Extremes of Heat and Cold was that of *De L'ISLE*, which was divided from the Point indicating the *Heat of boiling Water* 10600 Degrees, above and below ; and in some he mentions more than 800 to the

Bulb,

Bulb, which was, in his, of a *spherical* Form. But as *Fahrenbeit's* Scale is that in common Use, I have placed it by *De L'Isle's* on one Side, and that of *SIR ISAAC NEWTON* (as is generally supposed) on the other, as you observe in the Copper-plate of the *Grand Thermometer* annexed.

MR. BRAUNIGS observes, that neither *Fahrenbeit*, *Raumur*, *Muschentrock*, nor any other Person before himself, had been able to produce an artificial Cold which would cause the \varnothing to descend more than 40 Degrees below the Cypher (0) in *Fahrenbeit's* Scale, which was 110 in his own; but that taking the Advantage of a very cold Air at *Petersburgh* (*Dec* 14, 1759), the \varnothing standing at 205, he made a freezing Mixture with pounded Ice and *Aqua Fortis*, which, upon immersing the Bulb of the Thermometer, sunk the Mercury to 134 Degrees; which was 24 Degrees lower than had ever been seen before.

Flushed with this Success, he resolved to try what might be done with a Mixture of Snow and *Aqua Fortis*, and accordingly made one, in which, placing his Thermometer, the \varnothing sunk to 260, with a very few Drops of *Aqua Fortis*, and by putting in a little more, it sunk to 380 immediately.

MR. BRAUNIGS had ready by him another Glass filled with Snow, and taking the Thermometer out of the former, before it lost much of its Cold, he placed it in this second Mixture, and was agreeably surpris'd to observe the Mercury descend 470 Degrees

He says, such a stupendous Descent of \varnothing induced him to think, at first, that it could not be owing to cold, but to a Fracture of the Bulb of his Thermometer, which, when he had taken out, he was rejoic'd to find was entire and unhurt.

He now observed, for the first Time, that the \varnothing was no longer moveable in the Tube, but remained without Motion

STANDARD THERMOMETER.

3

Motion 12 Minutes in the open Air, cold to the Degree 203, and being carried into a Room 125 warm, after a few Minutes, the \varnothing recovered its Fluidity, and began to ascend

He now began to consider what might be done with other Sorts of Fluids in the Production of artificial Cold; with this View he tried various frigorific Mixtures, made with the following Liquors, and observed the Degrees through which the \varnothing descended by each, as in the Table below

Common Wines	- - - -	6, 7, 8.
Spirit of Sal. Ammoniac	- - - -	10.
Spirit of Sulphur	- - - -	10.
Spirit of Hart's Horn	- - - -	10.
Brandy	- - - -	12.
Spirit of Wine Camph.	- - - -	15.
Alcohol	- - - -	20.
Sweet Spirit of Vitriol	- - - -	20.
Spirit of Sea Salt	- - - -	30.
Simple Spirit of Nitre	- - - -	30.
Hoffman's Anodyne Liquor	- - - -	32.
Oil of Vitriol	- - - -	35.
Aqua Fortis	- - - -	40.
Fuming Spirit of Nitre	- - - -	58.

By these Experiments he was convinced, there was only one Liquor left for him to try, viz. the *Fuming Spirit of Nitre*, and this he did next use with amazing Success; for having provided several Glasses filled with Snow, on *December 25*, he poured the Spirit of Nitre in one, and the \varnothing subsided to 530; it there became motionless, and broke the Bulb of the Thermometer. He now first observed the \varnothing solid, though not totally so, for a small Quantity in the Middle of the Ball remained fluid. The external Surface of the mercurial Globe was continuous, polite, and hard; so that being put into a Mor-

tar, it was beaten flat with the Pestle, and appeared like Lead, emitting the same sound when struck; being reduced to thin Plates, they were cut with a Knife, and resembled the Brightness of the purest Silver, but in about twelve Minutes they began to melt, and soon recovered the usual Fluidity.

On *December 26*, the natural Cold of the Air exceeding any before, being at 212, Mr. *Braun* took the Opportunity of repeating his last Experiment in two Thermometers, and with such Success, that he observed the \varnothing to sink down to 650, and examining the Bulbs, found they were both cracked in several Parts, but yet adhered to the Mercury, which he observed was now just thoroughly congealed, and therefore he makes this Degree 650, the *Term of Cold for the Congelation or fixing of \varnothing into a solid Metal*.

By repeating these Experiments under various Circumstances, he was at length able to produce Descents of the \varnothing to 680 and 700 Degrees, and at last to 800, and beyond. In this prodigious Degree of Cold, the Bulb broke to Pieces, the \varnothing fell from the Height of three Feet to the Pavement, in Form of a solid Mass, or Globe of *Mercurial Metal*, it was flatted by the Fall; in short it was perfectly solid, ductile, malleable, &c in all respects like a Globe of Lead, Tin, or Silver.

Thus all Degrees of Cold to that which congeals and consolidates fluid Quicksilver into a hard Metal, have been explored and discovered by Experiment; and are express'd in all the three Scales affixed to this GRAND THERMOMETER, as you see in the Plate.

The upper Part of this Thermometer expresses all Degrees of Heat to that with which Bodies appear *glowing* in Twilight. Mr. *Braun* has found by his Experiments the Degrees of Heat with which Tin, Lead, Bismuth, &c. will melt, and Mercury boil; and these,
with

with the Heat of Melting Wax, Boiling Water, Summer Heat, &c are likewise marked on the Plate.

By Mr. *Braun's* Experiments it further Appears that those very Liquors which produced such remarkable Degrees of Cold by their Mixture with Snow, will at the same time create considerable Degrees of Heat if mixed with Water, even the Water of dissolved Snow. This Paradox in Philosophy requires too prolix a Solution for the Present, if it will admit of any at all.

Mr. *Braun's* Thermometer contains 1064 Divisions between boiling Mercury 414, and the Cold which congeals it to a Solid 650; and therefore the Bulk of the Mercury in its boiling State, being made equal to 10000 such Parts or Divisions, it is evident, that its Bulk in a State of Solidity will be 8936 of those Parts; and therefore it's Diminution in Bulk will be $\frac{1064}{10000}$, or nearly $\frac{1}{10}$ of the Whole; and of course it's Weight, or Specific Gravity, will be so much increased. But from a Temperate State of the Air to the Term of Congelation, there are but about 520 Divisions, therefore its Contraction in Bulk when solid, will be only $\frac{520}{10000}$, or about $\frac{1}{20}$ of its ordinary Bulk in a *fluid State*. It's Specific Gravity is now 14, but when made solid by Cold, it will be 14,7, and in its Boiling State, it will be 13,3.

Dr. *George Martine*, in his Treatise on the Thermometer, has given us a Plate of no less than 16 different Scales of Degrees or Divisions for making Observations with this Instrument, used by different Persons and Societies; among these only two begin the Scale at the *Freezing Point*, and Sir *Isaac Newton* was the first of these, and no wonder, since this great Man could do Nothing but what was consonant to the highest Reason, which undoubtedly points out that Degree of Temperature in the Air, by which the most common Fluids begin to freeze or congeal, as the proper Term of Comparison for
Heat

Heat and Cold, in all Degrees above or below it; if there be any Reason in the Divisions of the other Scales (particularly that of *Farenheit's* in common Use) I confess I am at a Loss to find it.

By the *Decimal Divisions* of the *Newtonian Scale* all Comparisons of Heat and Cold are greatly facilitated; thus if 12 represent the Heat of the Humane Body, then 24 the Heat of *melting Wax* is twice as great, and the Heat which *melts Tin* being 72 appears to be 6 Times as great, and, lastly, the Heat of *boiling Mercury* is 126 which is 10½ Times that of Blood, Milk, Urine, &c. On the other Hand, the Cold of a common *freezing Mixture*, here at *London*, is also 12, and the Cold producing the utmost Condensation of Mercury being 146, is more than 12 Times that of the Mixture.

But *Thermometers*, as they are commonly made, are little else than mere *Thermoscopes*, that is, they rather *show or indicate* the same relative Degrees of Heat and Cold, than *measure* them, as the Word *Thermometer* imports. Now in Order that they may be constructed to measure any Degrees in the Qualities of Heat and Cold, it must be considered that equal Degrees of Heat can only be measured by equal Expansions of the Fluid, and therefore the several Degrees on the Scales of Thermometers ought to be such as will adequately correspond to, and measure the equal Differences of those Expansions of the Fluid, above or below some given Point; and that in the most immediate and facile Manner.

To this End, it will be necessary to fix on some Point, as a *Term or Boundary* between Heat and Cold, And as *Sir Isaac Newton* has chosen this Term, which expresses that Temperament of the Air in which *Water begins to freeze, or Snow begins to thaw*, it would be presumption to think of any other for the BEGINNING OF THE SCALE

The

The Principle which Sir *Isaac Newton* and *De L'Isle* go upon is undoubtedly the best, but as OIL and SPIRIT are far from being proper Fluids for an universal Standard Thermometer, theirs could not be used in such a character; nor can any Fluid be found for this Purpose but the purest MERCURY, as appears from what has been said above.

The most convenient Form for the Bulb is by no means a *Globe*, as the Heat or Cold cannot so readily and instantly penetrate thro' such a Body, it ought to be therefore of a long and flatted Figure nearly like that in the Plate, and the Tube having a proper Length, and Size of the Bore, is first to be nicely weigh'd in a most exact Balance, and then to be filled with Mercury to the Top, as it stands in thawing Snow or Ice.

Secondly; The Bulb is then to be immersed into *boiling Water*, the Heat of which will expel a certain Quantity of \varnothing from the Tube, which must be carefully collected, and its Weight taken in Grains.

Thirdly, The length of the Bore left vacant by the Expulsion of the said \varnothing must be exactly measured in *Inches* 100th Parts.

Fourthly, This being done, the Bulb of the Thermometer is to be held over the Flame of a Spirit-Lamp till it has driven out so much \varnothing , that the remainder may stand about the middle of the Stem when the Bulb is again put into melting Snow, and thus you will have the *Freezing Point* ascertained, to be marked with a File.

Fifthly, From this Point you set off the before measured Distance or Length of \varnothing expell'd by boiling Water, and that will assign the *Point of Boiling Water* in the Scale.

Sixthly, The Thermometer as now fixed to the freezing Point is to be very exactly weighed, and from this Weight deduct that of the empty Tube, and the Remainder will be the Weight of the contained Mercury

Seventhly,

Seventhly ; Then say, As this whole Weight of φ to the freezing Point is to the Weight of that driven out by boiling Water, so is 10000 to a fourth Number A ; this Number A will therefore express how many of the 10000th Parts of the whole φ are contained in the Mercury between the Points of Freezing and Boiling Water.

Eighthly ; Then say, As the Number A is to the whole Distance between these two Points, so is 100 to the Length of the Bore occupied by 100th Part of the whole Bulk of $\varphi = 10000$.

Ninthly ; This Distance or Measure of 100 Parts is carefully to be set off from the Freezing Point above towards the Top of the Tube, and below towards the Bulb and each of these 100ths being divided into 10 equal Parts, they will adequately shew the Expansion of the Mercury in 10000th Parts of the Whole. And thus the Scale is completely finished for Use.

In all that has been hitherto said, it is supposed that the Bore of this Standard Thermometer-Tube is every where equal ; which ought to be examined at first by letting a small Quantity of φ move thro' the whole Length of the Tube, and then observing if it has the same Length in every Part ; if so, it is fit for the Purpose, otherwise not.

The Number A above-mentioned, I have found in several Trials to fall between 150 and 155 ; and particularly in two Tubes with flat Bulbs it came out 153 to the 8th Part of a Grain. I have therefore given 153 Parts of 10000 to the Distance between the Freezing Point, and that of Boiling Water, in the *Standard Thermometer* of the annexed Plate.

Whoever will consult Dr. G MARTINE'S Essays on these Subjects, will be satisfied that none of the numerous Constructions of this Instrument, which he has exhibited, are fitted for a Standard in this Kind. He has, with great
Judgment

Judgment, shewn, that the said Number A which *Fahrenheit* makes 180, *De L'Isle* 150, and *Reaumur* 80, cannot be depended upon for Exactness, but we have given them a Place here, as they serve to communicate and compare Observations made in different Parts of the World; as that of *Reaumur* in *France*, *De L'Isle's* at *Petersburgh*, and *Fahrenheit's* all over the Globe.

But as no Thermometer has yet been made to that Extent of Scale, as Nature and Use require, I have thought it might be acceptable to the curious and philosophic Part of the Public to have this first Exemplar of such a wonderful Construction. Dr. MARTINE was astonished at a Degree of Cold which caused a Descent of the \varnothing to 275 below (0) in *De L'Isle's* Thermometer at *Petersburgh*, he could scarce think it credible, but what would he have thought, or said, on hearing that in the Year 1760 it fell to 7 or 800 Degrees! and congealed the \varnothing , which the Dr. had denied to be possible! But what will not Time produce? We make such Thermometers now (for such as are willing to repeat those amazing Experiments in very cold Climates) with the utmost Ease, and though that Extreme of Cold cannot be exhibited here, yet nothing is easier than to shew the other Extreme of Heat for boiling \varnothing in the STANDARD THERMOMETER. And, by several Experiments, I have found it to be very near the same with that of Mr. *Braun*, as here marked in the Scales.

Philosophers have proposed a Method of investigating Degrees of Heat much greater than that of \varnothing , when it boils, even to that of *glowing*, or *red hot Metals*, and, of course, to explore it where no Thermometer can be applied. They go upon this Supposition, *that the Degrees of Heat lost by Bodies in equal Times, are as the remaining Degrees of Heat in those Bodies at the Ends of those equal Intervals of Time*; and if this were the Case, then the said Degrees of

C

lost and remaining Heat would be in *geometrical Proportion*, while the Intervals of Time were in *arithmetical Proportion*; and granting this, nothing could be easier, than, (by observing the Time in which the Heat of red hot Metal became equal to that which the Thermometer could bear when applied to it, and afterwards the Time (in Minutes) in which the \varnothing subsided to the Temperature of the ambient Air,) to find the Degree of Heat in the glowing Metal when first taken from the Fire; for, by this Means, the *logarithmic CURVE* might be easily drawn to express all the Degrees of Heat in the Mercury and in the Metal corresponding to every Minute of Time during the Observation, in the Manner directed in my *New and Compensatus SYSTEM of LOGARITHMS*.

But the Virtuoso has little Reason to hope for much Satisfaction in this Affair, for that the lost Degrees of Heat are as the Remainders in equal Times, is an *assumed Position*, rather than a *real Truth*; besides, were it actually Fact, the Heat of the \varnothing could never become equal to that of the common Air, as is evident from the Nature of the before-mentioned Curve and it's Asymptote; nor should I have troubled the Reader with such an *Hypothesis*, had it not been ascribed to SIR ISAAC NEWTON, and so largely treated of by Dr. G. MARTINE.

F I N I S.